



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hongkong.—During the period from the 10th to the 23d May there were registered 253 cases of plague, of which 222 ended in death.

British South Africa.—In Cape Colony there were registered during the week ended June 6, in Port Elizabeth, 9 new cases of plague (and 3 deaths); in King Williams Town, 5 cases (1 death). In East London on the 10th of June there occurred 1 case of plague. The plague hospital in Saldanha Bay has been closed.

In Natal, during the week ended May 23, 2 fresh cases of plague were registered. At the end of the week there were still 6 plague patients under treatment in the hospital. Up to the present date there has occurred a total of 205 plague cases and 150 deaths.

Plague and cholera.

British India.—In Calcutta during the week ended May 30 there were registered 52 deaths from plague and 52 deaths from cholera.

Death rate of Berlin.

The death rate of Berlin for the week ended June 27 amounted, calculated on the year, to 13.8 per thousand of the population, this rate being lower than that of two-thirds of the other large German towns. The following cities had a considerably higher death rate than Berlin, viz: Bremen, Kiel, Hamburg, Braunschweig, Magdeburg, Hanover, Frankfort-on-the-Main, Leipzig, Munich, Nuremberg, Carlsruhe, Cologne, Halle, Stettin, Konigsberg, Breslau, and also Paris. The following towns had a lower death rate than that of Berlin, namely: Stuttgart, Dresden, Dusseldorf, Kassel, Schoneberg (with 10.1), Charlottenburg (with 9.8 per thousand). There was a slight decrease in the number of deaths among children in the first year of life, the mortality rate among infants being 4.0 per year and mille, less than half the rate of Stettin and Halle and much lower than that of Munich, Cologne, and Leipzig. There was a slight decrease in the number of cases of acute intestinal disease and diseases of the respiratory organs, the former causing 47, the latter 38, deaths. There were 69 deaths from phthisis pulmonalis and 44 deaths from cancer. Further, there were registered 6 deaths from diphtheria, 4 deaths from scarlet fever, 12 deaths from measles, and 12 persons died by violence.

GUATEMALA.

Report from Livingston, fruit port.

Acting Assistant Surgeon Peters reports as follows: Week ended July 13, 1903. Present officially estimated population, 3,500; five deaths; prevailing diseases, malarial; general sanitary condition of this port and the surrounding country during the week, good.

Bills of health were issued to the following-named vessels:

Date.	Name of vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
July 7	David	19	0	0	0
7	Breakwater	34	2	0	2

The *Breakwater* cleared from Puerto Barrios.